



<https://www.biodiversitylibrary.org/>

**Prodromus systematis naturalis regni vegetabilis, sive,
Enumeratio contracta ordinum generum specierumque
plantarum huc usque cognitarum, juxta methodi naturalis,
normas digesta**

Parisii, Sumptibus Sociorum Treuttel et Wurtz, 1824-73

<https://www.biodiversitylibrary.org/bibliography/286>

v.15:pt.2 (1862-1866): <https://www.biodiversitylibrary.org/item/7166>

Page(s): Page 109

Holding Institution: Missouri Botanical Garden, Peter H. Raven Library

Sponsored by: Missouri Botanical Garden

Generated 25 March 2024 10:56 AM

<https://www.biodiversitylibrary.org/pdf4/1684820i00007166.pdf>

This page intentionally left blank.

foliis e basi sessili attenuatâ linear-lanceolatis vel linear-oblongis obtusis vel acutiusculis, umbellaribus abbreviatis, floralibus ovato-rotundis liberis obtusissimis involucro subbrevioribus, involucro hemisphærico-campanulato fuce cum lobis ovatis subtruncatis 2-3-dentatis hirti, glandulis 4-5-transverse oblongis truncatis vel obsolete bicornutis, stylis brevibus bifidis apice incrassatis, capsula subdepressæ trisulcæ coccis rotundatis, semine ovato superficialiter rugoso-scorbiculato, carunculâ depressâ. ♂ In Maderâ (Webb! Lemann! Mason!). Vulgo *Figueiro de Inferno*.—Jacq. Schönbr. IV, t. 485. Folia 1-2 pollicès longa, 3-4 lineas lata. Radii umbellæ pollicares. Capsula piso paulo minor. (v. s.)

424. E. REGIS JUBÆ (Webb phyt. Canar. III, p. 250), fruticosa, glabra, ternatim et quinatim ramosa, umbellæ 6-10-radiatæ, radiis bifidis, foliis e basi sessili attenuatâ anguste linearibus obtusis vel retusis, umbellaribus abbreviatis, floralibus ovatis obtusis involucro sublongioribus, involuci turbinati lobis ovato-triangularibus valde hirtis, glandulis truncato-semilunatis breviter et crasse bicornutis, stylis longis basi coalitis apice bifidis subincrassatis, capsulæ profunde trisulcatæ coccis rotundatis, semine ovato obsolete ruguloso-malleato, carunculâ ovato-cristæformi. ♂ In rupestribus Tenerifæ, Canariæ, Lancerottæ et Palmæ (Webb! Bourgeau n. 1513!). E. piscatoria Link in Buch Can. p. 158, non Ait. E. Mauritanica Webb florul. Lancerott. p. 31 olim, non L. A præcedenti foliis angustioribus (2 lineas latis), glandulis evidentius bicornutis, stylis duplo longioribus, semine obsolete scrobiculato distincta. (v. s.)

425. E. BERTHELOTI (C. Bolle in litt.), arborescens, glabra, truncto brevi basi crassissimo transverse et annulatim incrassato, ramis cinereis apice incrassatis rubescientibus comam latissimam laxam formantibus post anthesin aphyllis, foliis sessilibus late linearibus basi et apice attenuatis obtusissimis apice retusis vel mucronulatis glaucescenti-viridibus subtus pallidioribus, umbellæ radiis 6-10 bifidis, involuero.... ♂ In regione calidâ littorali insulæ Canariensis Gomera, loco Barranco de San Sebastian dicto (Bolle). Arbuscula 5-7-pedalis, ramis crassis angulo obtusissimo dichotomis comam diametro 5-6-pedalem formantibus. Folia 2 pollices longa, 5-6 lineas lata, autumno decidua, nec ut in affinibus in rosulam persistentia. Inflorescentia eam E. Regis Jubæ referens. Hanc speciem adhuc imperfecte notam ex auctoritate cl. Bolle descripti qui eam habitu insignem et ab affinibus distinctissimam esse dicit.

426. E. TUCKEYANA (Steud. nom. I, p. 615), glabra, fruticosa vel arborescens, ramis inferne denudatis superne densiuseule foliosis, umbellæ 4-5-radiatæ radiis bifidis, foliis sessilibus linearibus obtusis basi sensim attenuatis, umbellaribus oblongis radiis paulo brevioribus, floralibus ovatis obtusis involucro vix longioribus, involuci campanulati glabri intus pubescentis lobis acute bidentatis, glandulis ovatis margine truncato-retusis vel brevissime bicornutis, stylis fere ad medium coalitis breviter bifidis apice non incrassatis, capsulâ globoso-subtriquetrà sub lente elevato-punctatâ coccis obsolete carinatis, semine ovato lævi fusco-marmorato carunculâ pileatâ. ♂ In altioribus insularum San Vincent et San Antonio archipelagi Capitis Viridis (Tuckey! Smith!), in insulâ San Nicolas (Bolle!) E. arborea Chr. Smith in Tuckey Voy. p. 251. Schmidt Fl. Cap. Verd. p. 304. Frutex vel arbuscula 6-8 pedalis, basi 3-4 poll. diam. lata. Folia 2-3 poll. longa, 3 lin. lata. Radii 2-2½-pollicares. Capsula piso quartâ parte minor. Ab affini E. Regis Jubæ differt foliis non retusis, umbellâ pauciradiatâ, foliis floralibus majoribus, lobis involuci bidentatis et semine lævi. (v. s. comm. a cl. Dr. Bolle et in h. Kew.)

427. E. DENDROIDES (Linn. sp. p. 662), fruticosa, glabra, dichotome vel ternatim ramosa, ramis rubellis nitidis annotinis dense foliosis, umbellæ 3-10-fidæ radiis brevibus bifidis, foliis sessilibus e basi valde attenuatâ linear-lanceolatis obtusis vel acutiusculis subtus pallidioribus, umbellaribus abbreviatis, floralibus lutescentibus involucrum superantibus subcordato vel rhombo-semiorbiculatis mucronulatis vel muticis, involuci campanulati lobis oblongis 2-3-fidis fimbriato-ciliatis, glandulis peltatis transverse ovatis subtus hirtulis truncatis vel semilunatis, stylis elongatis tenuibus ad tertiam partem connatis bifidis apice subincrassatis, capsulæ profunde trisulcæ coccis lateraliter subcompressis, semine ovato lævi subpresso, carunculâ sessili oblongâ cristæformi. ♂ In rupestribus calidioribus maritimis regionis Mediterraneæ in Catalauniâ (h. DC!), Galliâ australi! Italiâ et Græciâ!, Cretâ (Sieber!), Afrieâ boreali! — Sibth. fl. Græc.

The following text is generated from uncorrected OCR or manual transcriptions.

[Begin Page: Page 109]

EUPHORBIA. EUPHORBIEAZE (vcro BOISSIER]. 109

foliis e basi sessili attenuatà linear-i-lauceolatis vel linear-i-oblongis obtusis vel acutiusculis, umbellaribus abbreviatis, floralibus ovato-rotundis liberis obtusis-simis involuero subbrevioribus, involucro hemisph:erico-campanulato fauce cuni lobis ovatis subtruncatis 2-3-dentatis hirto, glandulis 4 5-transverse oblongis truncatis vel obsolete bicornutis, stylis brevibus bifidis apice incrassatis, capsulie subdepresso trisuleze coccis rotundatis, semine ovato superficialiter rugoso-scrobiculato, carunculà depressá. 5 In Maderà (Webb! Lemann! Mason !). Vulgo Fi-gueiro de Infrerno.— Jacq. Schónbr. IV, t. 485. Folia 1-2 pollices longa , 3-4 lineas lata. Radii umbellie pollicares. Capsula piso paulo minor. (v. s.)

424. E. Reis Jun & (Webb phyt. Canar. III, p. 250), fruticosa, glabra, ternatini et quinalim ràmosa, unibell:e 6-10-radiatze, radiis bifidis, foliis e basi sessili atte-nuatà anguste linearibus obtusis vel retusis, umbellaribus abbreviatis, floralibus ovatis obtusis involucro sublongioribus, involuci turbinati lobis ovato-triangu-laribus valde hirtis, glandulis truncato-semilunatis breviter et crasse bicornuti»s, stylis longis basi coalitis apice bifidis subincrassatis, capsul: profunde trisulcat-coccis rotundatis, semine ovato obsolete ruguloso-malleato, carunculà ovato-cristeformi. 5 In rupestribus Tenerife, Canarie, Lancerotiz et Palm (Webb! Bourgeau n. 15131). E. piscatoria Link in Buch Can. p. 158, non Ait. E. Maurita-niea Webb florul. Lancerott. p. 31 olim, non L. A pr:ecedenti foliis angustioribus (2 lineas latis), glandulis evidentius hicornutis, stylis duplo longioribus, semine obsolete scrobiculato distincta. (v. s.)

425. E. BER THELOT I (C. Bolle in lit.), arborescens, glabra, trunco brevi basi cras-sissimo transverse et annulatim incrassato, ramis cinereis apice incrassatis ru-bescientibus comam latissimam laxam formantibus post anthesin aphyllis, folii sessilibus late linearibus basi et apice attenuatis obtusissimis apice retusis vcl mucronulatis glaucescenti-viridibus subtus pallidioribus, umbell:& radiis 6-10 bifidis, involucro... 5 In regione calidà littorali insulze Canariensis Goméra, loco Barranco de San Sebastian dicto (Bolle). Arbuscula 5-7-pedalis, ramis crassis ungulo obtusissimo dichotomis comam diametro 5-6-pedalem formantibus. Folia 2 pollices longa, 5-6 lineas lata, autumno decidua, nec ut in affinibus in rosulazi persistentia. Inflorescentia eam E. Regis Juba referens. Hanc speciem adhuc im-perfecte notam ex auctoritate cl. Bolle descripsi qui eam habitu insignem et ab affinibus distinctissimam esse dicit:

14126. E. TUcKEYANA (Sleud. nom. I, p. 615), glabra, frutiecosa vel arborescens, ramis inferne denudatis superne densiuscule foliosis, umbella 4j-5-radiati radiis bifidis, foliis sessilibus linearibus obtusis basi sensim attenuatis, umbellaribus oblongis radiis paulo brevioribus, floralibus ovatis obtusis involucro vix longio-ribus, involuci campanulati glabri intus pubescentis lobis acute bidentatis, glan-dulis ovatis margine truncato-retusis vel brevissime bicornutis, stylis fere ad medium coalitis breviter bifidis apice non incrassatis, capsulá globoso-subtri-quetrà sub lente elevato-punctatà coccis obsolete carinatis, semine ovato levi fusco-marmorato carunculá pileatà. 5 In altioribus insularum San Vincent et San Antonio archipelagi Capilis Viridis (Tuckey | Smith I), in insulá San Nicolas (Bolle !) E. arborea Chr. Smith in Tuckey Voy. p. 251. Schmidt Fl. Cap. Verd. p. 304.

Frutex vel arbuscula 6-8 pedalis, basi 3-4 poll. diam. lata. Folia 2-3 poll. longa, 3 lin. lata. Radii 2-22-pollicares. Capsula piso quartà pártē minor. Ab affini E. Regis Jub: differt foliis non retusis, umbellà pauciradiatà, foliis floralibus majo-ribus, lobis involuci bidentatis et semine levi. (v. s. comm. a cl. Dr. Bolle et inh.Kew.) |

497. E. pENDnoOIDES (Linn. sp. p. 662), fruticosa, glabra, dichotome vel terna-

tim ramosa, ramis rubellis nitidis annotinis dense foliosis, umbellz 3-10-fida* radiis brevibus bifidis, foliis sessilibus e basi valde attenuatà linear-i-lanceolatis obtusis vel aeutiusculis subtus pallidioribus, umbellaribus abbreviatis, floralibus lutescentibus involucrum superantibus subcordato vel rhombeo-semiorbiculatis mucronulatis vel mulicis, involucri campanulati lobis oblongis 3-3-fidis firmato-ciliatis, glandulis peltatis transverse. ovatis subtus hirtulis truncatis vel semilunatis, stylis elongatis tenuibus ad tertiam partem connatis bifidis apice subinerassatis, capsule profunde trisulez coccis lateraliter subcompressis, semine ovato levi subpresso, carunculà sessili oblongà crist:iformi. 5 In rupestribus calidioribus maritimis regionis Mediterranee in Cataleunià (h. DC), Gallià ' australi! Italià et Greecià !, Cretà (Sieber!), Afrieà borcali!— Sibth. il. Grec.