



<https://www.biodiversitylibrary.org/>

**Hortus Kewensis, or, A catalogue of the plants cultivated  
in the Royal Botanic Garden at Kew**

London, Printed for George Nicol, Bookseller to his Majesty, 1789  
<https://www.biodiversitylibrary.org/bibliography/4504>

**v.3:** <https://www.biodiversitylibrary.org/item/23434>

Page(s): Page 107

Holding Institution: Missouri Botanical Garden, Peter H. Raven Library  
Sponsored by: Missouri Botanical Garden

Generated 29 March 2024 12:02 PM  
<https://www.biodiversitylibrary.org/pdf4/1686203i00023434.pdf>

This page intentionally left blank.

Trailing St. John's-wort.

Nat. of Britain.

Fl. July.

H. 4.

19. H. floribus trigynis, calycibus ferrato-glandulosis, caule tereti erecto, foliis ovatis glabris. *Syst. veget.* 701. *montanum.*

Mountain St. John's-wort.

Nat. of Britain.

Fl. July.

H. 4.

20. H. floribus trigynis, calycibus acutis ferrato-glandulosis, foliis oblongo-lanceolatis pellucido-punctatis margine glandulosis, caule fruticoso. *glandulosum.*

Glandulous St. John's-wort.

Nat. of Madeira. Mr. Francis Masson.

Introd. 1777.

Fl. May—August.

G. H. 5.

21. H. floribus trigynis, calycibus ferrato-glandulosis, caule tereti erecto, foliis ovatis subpubescentibus. *hirsutum.*

*Sp. pl.* 1105. *Curtis lond.*

Hairy St. John's-wort.

Nat. of Britain.

Fl. July.

H. 4.

22. H. floribus trigynis, calycibus ferrato-glandulosis, foliis femiamplexicaulibus flexuosis tomentosis, caulibus prostratis. *Sp. pl.* 1106. *tomentosum.*

Woolly St. John's-wort.

Nat. of the South of Europe.

Introd. 1772, by Mons. Richard.

Fl. July—September.

G. H. 4.

23. H. floribus trigynis, caule subancipiti, foliis amplexicaulibus ovatis, cyma floribus sessilibus. *Syst. veget.* 702. *perfoliatum.*

702.

Perfoliate

The following text is generated from uncorrected OCR or manual transcriptions.

[Begin Page: Page 107]

POLYADELPHIA POLYANDRIA. Hypericum.

Trailing St. John's-wort.

Nat. of Britain. | Soy

fi, July. . H, 4.

19. H. floribus trigynis, calycibus ferrato-glandulofis, caule

tereti erecto, foliis ovatis glabris.. Sy/f. wagels 104s

'Mountain St. John's-wort,

Nat. of Britain. ae

Fl, July. y ; H. UY.

20. H. floribus trigynis, calycibus acutis ferrato-glandu-

lofis, foliis oblongo-lanceolatis pellucido-punctatis

marginis glandulofis, caule fruticifera.

Glandulous St. John's-wort. \_

Nat. of Madeira. Mr. Francis Maffin.

Introd. 1777 E>

Fi. May—August. Bs: 9: Sas

21. H. floribus trigynis, calycibus ferrato-glandulofis,

\_caule tereti erecto, foliis ovatis fubpubefcentibus.

Sp. pl. 1105. Curtis lond.

Hairy St. John's-wort.

Nat. of Britain." ha

et faly- H. y.

22. H. floribus trigynis, calycibus ferrato-glandulofis,

foliis . femiamplexicaulibus fexuofis.. taencntohie,

caulibus profratis. Sp, pl. 11206.

Woolly St. John's-wort.

"Nat. of the South of Europe.

- Introd. 1772, by Monf. Richard.

Fi. July —September. ae G. H. Ue

23, HL. floribus. Sinznin peers fubancipiti, foliis amplexi-

caulibus ovatis, cyma floribus feffilibus, oe saa

702. ¢

ee

ies Pertoliate. 3 reas

107

moutaa-

RUM

glandulo- : :

fue

hirfutume

aes at a