



<https://www.biodiversitylibrary.org/>

**Caroli Linnaei ... Species plantarum :exhibentes plantas rite cognitatas, ad genera relatas, cum differentiis specificis, nominibus trivialibus, synonymis selectis, locis natalibus, secundum systema sexuale digestas...**

Holmiae, Impensis Laurentii Salvii, 1753

<https://www.biodiversitylibrary.org/bibliography/669>

**vol. 2:** <https://www.biodiversitylibrary.org/item/13830>

Page(s): Title Page, Page 1196

Holding Institution: Missouri Botanical Garden, Peter H. Raven Library

Sponsored by: Missouri Botanical Garden

Generated 5 January 2025 3:12 AM

<https://www.biodiversitylibrary.org/pdf4/1763774i00013830.pdf>

This page intentionally left blank.

CAROLI LINNÆI

S:Æ R:GIÆ M:ITIS SVECIÆ ARCHIATRI; MEDIC. & BOTAN.  
PROFESS. UPSAL; EQUITIS DE STELLA POLARI;

SPECIES  
PLANTARUM,

EXHIBENTES

PLANTAS RITE COGNITAS,

AD

*GENERA RELATAS,*

CUM

DIFFERENTIIS SPECIFICIS,  
NOMINIBUS TRIVIALIBUS,  
SYNONYMIS SELECTIS,  
LOCIS NATALIBUS,

SECUNDUM

*SYSTEMA SEXUALE*

DIGESTAS.

TOMUS II.

---

*Cum Privilegio S. R. Mitis Sueciæ & S. R. Mitis Poloniæ ac Electoris Saxon.*

---

HOLMIÆ,

IMPENSIS LAURENTII SALVII.

1753.

*arenaria.* 8. STELLARIA caule erecto bifido, ramis alternis, petalis emarginatis. (423. post 7.)

*Habitat in Hispania.*

Radix fibrosa, annua. Caulis spithameus, erectus, teres, pilis patulis pubescens & subglutinosus. Rami alterni, longitudine fere caulis. Folia opposita, sessilia, subtus & margine pilosa, supra vero vix manifeste, ligulata apice ovato, basin versus angustiora linearia. Primus flos e dichotomia caulis, ubi caulis bifidus evadit; reliqui ex alis ramorum alterni, omnes majusculi. Pedunculus erectus, foliis longior. Calyx pentaphyllus, viscosus uti tota planta, apicibus purpurascens uti etiam folia floralia. Corolla campanulata, calyce longior, obtusissima vel vix manifeste emarginata aut retusa, alba. Stamina alba, longitudine calycis. Antheræ cerulescentes. Germen globosum. Styli 3, longitudine staminum.

*hispanica.* 7. COTYLEDON foliis oblongis subteretibus, floribus fasciculatis. (430. post 6)

Cotyledon maritima, sedi folio, flore carneo, fibrosa radice. *Tournef. inst. 90.*

Cotyledon hispanica, sedi teretifolii folio, flore umbellato rubro, fibrosa radice. *Minuart. sched. 3.*

*Habitat in Hispania. Læfling. ♂*

Radix fibrosa. Caulis simplex digitalis, teres, erectiusculus. Folia oblonga, teretiuscula, obtusiuscula, sessilia, subpilosa, supra planiuscula, ruffo-punctata. Corymbus terminalis. Flores infundibuliformes. Corollæ erectæ, extus rufescentes, subpilosæ: limbo plano, quinquepartito, acuto, supra incarnato striis tribus purpurascens & apice purpureo. Antheræ minima, filamenta crassiuscula, prominula. Calyx brevis. Facies Herbe Sedi albi.

*dubium.* 9. PAPAVER capsulis oblongis glabris, caule multifloro fetis adpressis foliis pinnatifidis incis. (507. post 5.)

Argemone capitulo longiore glabro. *Moris. bles. 233. hist. 2. p. 279. f. 3. t. 14. f. 11.*

*Habitat inter Sueciæ, Angliæ segetes. ☉*

Planta media inter P. argemone & Rhæas. Capsulis clavatis & pilis pedunculo adpressis cum P. Argemone convenit & a P. Rhæde differt.

*squamatus.* 29. CISTUS suffruticosus stipulatus, foliis obtectis squamis orbiculatis (526. post. 13.)

*Habi-*

The following text is generated from uncorrected OCR or manual transcriptions.

[Begin Page: Title Page]

> CAROLI LINNÆI

S: gŕ R: cœ M: ris Sveciœ ARculAaTRI; Mapic. & Bora,  
PRorrss. UrsAr; EQUITIS DE STELLA POLARI;

SPECIES

PLANTARUM,

EXHIBENTES

PLANTAS RITE COGNITAS,

AD

GENERA RELATAS,

CUM

DIFFERENTIIS SPECIFICIS,

NoMiNIBUs 'TRIVIALIBUS,

SYNONYMIS SELECTIS,

Loci NaTALiBUs,

t

SECUNDUM

SYSTEMA SEXUALE

DIGESTAS.

Towuusll.

Cum Privilegio S, R. Mitis Suecia. id. S. R. Mtis Polonie ac Elettoris Saxon,

T S TT TU MONT NIIS IS TUIS T DS ND ee

HOLMIZ,

'IMPENSIS LAURENTII SALVIL

17535,

[Begin Page: Page 1196]

rt9ó APPENDIX.

atiis ^ 8. STELLARIA canle ere&o bifido, ramis alternis, pe"  
talis emarginatis. (423. poft 7.)

Habitat in Hifpania.

Radix fibrofa; annua. Caulis fpitbameus, erectus, teres,

pilis patulis pubefcens y [ubglutino[us.. Rami alterni,

longitudine fere caulis. Y olia oppofita , fefilia fub-

ſus C9 margine pilofa, [upra vero vix manifeſte, li-

ulata apice ovato, bafim verſus anguſtiora linearia.

Érimus flos e dichbotomia caulis, ubi caulis bifidus e-

~ vadit; reliqui ex alis rdmorum alterni; omucſtomajn-

feculi. Pedunculus eregas , foliis longior. Calyx pen-

zaphyllus , vifcoſus uti tota planta, apicibus purpura-

feentibus uti etjam folia. floralia. Corolla campanula-

za, talyce. longior , obtufiffima vel vix manifeſte e7

- marginata aut retufa, alba. Stamina alba, longitudi-

ne calycis. Antherg ceruleſcentes, Germen globofume.

lia, fubpilofa, Jafra laniufcula, ruffo-punétata. Go-

rymbus žerminalis. Flores infandibuliformes, Coto-

purpurafcentibus E apice purpureo. Antherg mini-

Pe, Filamenta ENAA , prominula. "Calyx bre-

vis, Facies Herbe Sedi albi. :

Sm, 9, PAPAVER capfulis oblongis glabris, caule multiflo-  
ro fetis adpreiis foliis pinnatifidis incifis. (507.p9/? 5-)

Argemone capitulo longiore glabro. Mor:f.-olef. à33-  
iff. 2. p. 279. f 3. ti 14. f. 1T. i À

Habitat inter Suecie, Anglie fegetes: "© > FH

Planta media inter' P. argemoüe €9" Rheas. Capfulis  
elavatis £9» pilis pedunculo adpreffis cum P. Atge-  
mone cozvesuit & a P. Rhzde Ziffert. ^ =^

. A 1 , F Wt 2r 5 F,

fgudmatus, 29, CISTUS fuffruticofus flipulatus, foliis obtetis Íqua-  
mis orbiculatis (526. poft. 13.). Habi-